

OPEN  ACCESS



# International Journal of Applied Sciences and Biotechnology

A Rapid Publishing Journal

**ISSN: 2091-2609**

## Indexing and Abstracting

CrossRef, Google Scholar, Global Impact Factor, Genamics, Index Copernicus, Directory of Open Access Journals, WorldCat, Electronic Journals Library (EZB), Universitätsbibliothek Leipzig, Hamburg University, UTS (University of Technology, Sydney): Library, International Society of Universal Research in Sciences (EyeSource), Journal Seeker, WZB, Socolar, BioRes, Indian Science, Jadoun Science, Jour-Informatics, Journal Directory, JournalTOCs, Academic Journals Database, Journal Quality Evaluation Report, PDOAJ, Science Central, Journal Impact Factor, NewJour, Open Science Directory, Directory of Research Journals Indexing, Open Access Library, International Impact Factor Services, SciSeek, Cabell's Directories, Scientific Indexing Services, CiteFactor, UniSA Library, InfoBase Index, Infomine, Getinfo, Open Academic Journals Index, HINARI, etc.

**CODEN (Chemical Abstract Services, USA): IJASKD**

**Vol-4, Issue-1 (March, 2016)**



Impact factor\*: 1.422  
Scientific Journal Impact factor#: 3.419  
Index Copernicus Value: 6.02  
IBI Factor 2015\*\*: 4.19

Available online at:

<http://www.ijasbt.org>  
&  
<http://www.nepjol.info/index.php/IJASBT/index>



\*Impact factor is issued by Universal Impact Factor. Kindly note that this is not the IF of Journal Citation Report (JCR).

#Impact factor is issued by SJIF INNO SPACE; \*\*Impact factor is issued by INFOBASE INDEX.



## Research Article

# IN VITRO CALLUS PROLIFERATION FROM LEAF EXPLANTS OF GREEN GRAM AFTER *IN SITU* UV-B EXPOSURE

K Rajendiran\*, S Vidya and D Arulmozhi

Department of Botany, K.M. Centre for Post Graduate Studies, Pondicherry - 605 008, India.

\* Corresponding author's email: rajeworks@yahoo.com

### Abstract

Callus induction was tried with leaf explants (third leaf from top of canopy) harvested from *in situ* control and supplementary UV-B irradiated (UV-B = 2 hours daily @ 12.2 kJ m<sup>-2</sup> d<sup>-1</sup>; ambient = 10 kJ m<sup>-2</sup> d<sup>-1</sup>) three varieties of green gram *viz.* CO-8, NVL-585 and VAMBAN-2 to study their viability for germplasm conservation. Callus induction occurred both in control and UV-B stressed NVL-585 leaf explants. VAMBAN-2 both in unstressed and UV-B stressed conditions did not initiate callus. Only control leaf explants from CO-8 proliferated callus. Callus of UV-B irradiated NVL-585 weighed less (51.28 %) than control. Parenchyma cells were smaller in callus induced from *in situ* UV-B exposed NVL-585 leaf explants. The leaf explants from UV-B stressed NVL-585 varieties of green gram responded to *in vitro* callus proliferation making them fit for germplasm conservation for cultivating in UV-B elevated environment.

**Key words:** Callus proliferation; Green gram; Leaf explants; Three varieties; Ultraviolet-B.

### Introduction

Ozone depletion is facilitated by increased stratospheric particles as provided by stratospheric clouds in the polar regions and globally by volcanic eruptions. A drop in the level of ozone allows ultraviolet-B (UV-B) radiation (280-320 nm) into the earth surface creating a dangerous atmospheric stress to wild plants and domesticated crops (Caldwell *et al.*, 1998) as it destroys the leaf epidermis (Kokilavani and Rajendiran, 2013; Kokilavani and Rajendiran, 2014a; Kokilavani and Rajendiran, 2014b; Kokilavani and Rajendiran, 2014c; Kokilavani and Rajendiran, 2014d; Kokilavani and Rajendiran, 2014f; Kokilavani and Rajendiran, 2014g; Kokilavani and Rajendiran, 2014h; Kokilavani and Rajendiran, 2014j; Kokilavani and Rajendiran, 2014k; Kokilavani and Rajendiran, 2014l; Kokilavani and Rajendiran, 2014m; Kokilavani and Rajendiran, 2014n; Kokilavani and Rajendiran, 2015a; Kokilavani and Rajendiran, 2015b) even when plants in reflex pump epicuticular wax to leaf surface rapidly (Gowsalya and Rajendiran, 2015; Thiruvarasan and Rajendiran, 2015; Vidya and Rajendiran 2015) causes stomatal aberrations in cotyledonary epidermis of germinating seedlings (Rajendiran *et al.*, 2015c; Rajendiran *et al.*, 2015d) suppresses photosynthesis (Kulandaivelu *et al.*, 1989; Sullivan *et al.*, 1994; Rajendiran, 2001) causes abnormalities in morphology thereby reducing plant growth (Rajendiran and

Ramanujam, 2003; Rajendiran and Ramanujam, 2004; Kokilavani and Rajendiran, 2014o; Rajendiran *et al.*, 2015b; Rajendiran *et al.*, 2015k; Rajendiran *et al.*, 2015n; Rajendiran *et al.*, 2015p; Rajendiran *et al.*, 2015q), reduces yield (Mark and Tevini, 1997; Rajendiran and Ramanujam, 2004; Kokilavani and Rajendiran, 2014e; Rajendiran *et al.*, 2015m; Rajendiran *et al.*, 2015p; Rajendiran *et al.*, 2015q) and disturbs nodulation and symbiotic nitrogen fixation (Rajendiran and Ramanujam, 2006; Sudaroli Sudha and Rajendiran, 2013a; Sudaroli Sudha and Rajendiran, 2013b; Kokilavani and Rajendiran, 2014i; Sudaroli Sudha and Rajendiran, 2014a; Sudaroli Sudha and Rajendiran, 2014b; Sudaroli Sudha and Rajendiran, 2014c; Arulmozhi and Rajendiran, 2014a; Arulmozhi and Rajendiran, 2014b, Arulmozhi and Rajendiran 2014c, Vijayalakshmi and Rajendiran 2014a, Vijayalakshmi and Rajendiran, 2014b; Vijayalakshmi, Rajendiran 2014c; Rajendiran *et al.*, 2015l; Rajendiran *et al.*, 2015o) in many sensitive crops. Hence, *in vitro* screening needs to be carried out for selecting the germplasm of UV-B tolerant crop varieties prior to their conservation and regeneration. This study deals with identifying the varieties of green gram that can survive under supplementary UV-B irradiation and the possibilities of utilizing the leaf explants for germplasm preservation.

## Materials and Methods

### *In situ UV-B radiation*

Green gram (*Vigna radiata* (L.) Wilczek.), the nitrogen fixing grain legume was chosen for the study. Viable seeds of the three varieties of green gram viz. CO-8, NVL-585 and VAMBAN-2 were procured from Saravana Farms, Villupuram, Tamil Nadu and from local farmers in Pondicherry. The seeds were selected for uniform colour, size and weight and used in the experiments. The crops were grown in pot culture in the naturally lit greenhouse (day temperature maximum  $38 \pm 2$  °C, night temperature minimum  $18 \pm 2$  °C, relative humidity  $60 \pm 5$  %, maximum irradiance (PAR)  $1400 \mu\text{mol m}^{-2} \text{s}^{-1}$ , photoperiod 12 to 14 h). Supplementary UV-B radiation was provided in UV garden by three UV-B lamps (*Philips TL20W/12 Sunlamps*, The Netherlands), which were suspended horizontally and wrapped with cellulose diacetate filters (0.076 mm) to filter UV-C radiation ( $< 280$  nm). UV-B exposure was given for 2 h daily from 10:00 to 11:00 and 15:00 to 16:00 starting from the 5 DAS (days after seed germination). Plants received a biologically effective UV-B dose (UV-B<sub>BE</sub>) of  $12.2 \text{ kJ m}^{-2} \text{ d}^{-1}$  equivalent to a simulated 20 % ozone depletion at Pondicherry ( $12^{\circ}2'N$ , India). The control plants, grown under natural solar radiation, received UV-B<sub>BE</sub>  $10 \text{ kJ m}^{-2} \text{ d}^{-1}$ . Leaf explants (third leaf from top of canopy) were harvested from 30 DAS crops that received supplementary UV-B irradiation and sunlight in the *in situ* condition.

### *In vitro culture with leaf explants*

Leaf explants after appropriate aseptic treatment were used for *in vitro* culture. Leaf discs were thoroughly washed with water containing 0.1% Bavistin (a systemic fungicide BASF, India Ltd., Bombay) for 4-5 minutes. They were surface sterilized with 0.1% HgCl<sub>2</sub> for 4-5 minutes and washed 6 to 8 times with autoclaved water under Laminar Air Flow Cabinet (Technico Systems, Chennai) and inoculated aseptically onto culture medium. The final wash was given with aqueous sterilized solution of (0.1%) ascorbic acid. The surface sterilized explants were dipped in 90% ethanol for a short period (40 seconds).

The leaf discs were inoculated horizontally on MS medium for culture initiation. Different concentration and combination of cytokinins (6-benzyl amino purine – BAP and Kinetin ranging from 0.1 to 5.0 mg L<sup>-1</sup>) and auxins (IAA - Indole acetic acid ranging from 0.1 to 1.0 mg L<sup>-1</sup>) were incorporated in the medium for inducing callus. These cultures were incubated at  $28 \pm 2$  °C in the dark for 2-3 days. Subsequently these were kept under diffused light ( $22 \mu\text{mol m}^{-2} \text{s}^{-1}$  SFP- spectral flux photon) for 8 to 10 days. The light was provided by fluorescent tubes and incandescent bulbs. Temperature was maintained by window air conditioners. Positive air pressure was maintained in the culture rooms, in order to regulate temperature and to maintain aseptic conditions. The

cultures were regularly monitored and the callus proliferation were recorded after 30 DAI (days after inoculation). The experiments were carried out with three replicates per treatment.

The plant tissue culture media generally comprise of inorganic salts, organic compounds, vitamins, gelling agents like agar-agar. All the components were dissolved in distilled water except growth regulators. Auxins were dissolved in 0.5N NaOH or ethanol and cytokinins were dissolved in dilute 0.1N HCl or NaOH. For the present study MS basal medium (Murashige and Skoog 1962) was used as nutrient medium.

MS basal medium was used either as such or with certain modification in their composition. Sucrose and sugar cubes were added as a source of carbohydrate. The pH of the media was adjusted to  $5.8 \pm 2$  with 0.5N NaOH or 0.1N HCl before autoclaving. The medium was poured in the culture vessels. Finally the medium was steam sterilized by autoclaving at 15 psi (pounds per square inch) pressure at 121 °C for 15 minutes.

### *Chemical composition of MS medium (Murashige and Skoog 1962)*

Constituents	Quantity (mgL <sup>-1</sup> )
<b>Macronutrients</b>	
NH <sub>4</sub> NO <sub>3</sub>	1650
KNO <sub>3</sub>	1900
CaCl <sub>2</sub> .2H <sub>2</sub> O	440
MgSO <sub>4</sub> .7H <sub>2</sub> O	370
KH <sub>2</sub> PO <sub>4</sub>	170
Na.EDTA	37.23
FeSO <sub>4</sub> .7H <sub>2</sub> O	27.95
<b>Micronutrients</b>	
KI	0.83
H <sub>3</sub> BO <sub>3</sub>	6.20
MnSO <sub>4</sub> .4H <sub>2</sub> O	22.30
ZnSO <sub>4</sub> .7H <sub>2</sub> O	8.60
Na <sub>2</sub> MoO <sub>4</sub> .2H <sub>2</sub> O	0.25
CuSO <sub>4</sub> .5H <sub>2</sub> O	0.025
CoCl <sub>2</sub> .6H <sub>2</sub> O	0.025
Meso-Inositol	100
Glycine	2.0
Thiamine. HCl	0.1
Nicotinic acid	0.5
Pyridoxine. HCl	0.5
Sucrose (% w/v)	3 %
pH	5.8

### Preparation of MS medium

Approximately 90 % of the required volume of the deionized-distilled water was measured in a container of double the size of the required volume. Dehydrated medium was added into the water and stirred to dissolve the medium completely. The solution was gently heated to bring the powder into solution. Desired heat stable supplements were added to the medium solution. Deionized-distilled water was added to the medium solution to obtain the final required volume. The pH was adjusted to required level with NaOH or HCl. The medium was finally dispensed into culture vessels. The medium was sterilized by autoclaving at 15 psi at 121°C for appropriate period of time.

### Photography

The anatomical features were viewed through Nikon Labomed microscope under incident and translucent light and photographed using Sony digital camera fitted with Olympus adaptor. The culture tubes with leaf explants and callus were photographed in daylight using a Sony digital camera fitted with appropriate close-up accessories.

### Dendrogram

At least three replicates were maintained for all treatments and control. The experiments were repeated to confirm the trends. The result of single linkage clustering (Maskay 1998) was displayed graphically in the form of a diagram called dendrogram (Everstt 1985). The similarity indices between the three varieties of green gram under study were calculated using the formula given by Bhat and Kudesia (2011).

$$\text{Similarity Index} = \frac{\text{Total number of similar characters}}{\text{Total number of characters studied}} \times 100$$

Based on the similarity indices between the three varieties of green gram, dendrogram was drawn to derive the interrelationship between them and presented in Table 2 and Plate 5.

## Results and Discussion

### Standardisation of culture medium for leaf explant

For the standardisation of culture media leaf explants from green gram variety viz., NVL-585 grown under control condition were used (Plate 1). The explants were inoculated on MS medium for culture initiation containing different concentration and combination of cytokinins (6-benzyl amino purine - BAP = 2.0 mgL<sup>-1</sup> and Kinetin = 0.1, 0.25 and 0.5 mgL<sup>-1</sup>) and auxins (IAA - Indole acetic acid = 1.0 mgL<sup>-1</sup>). The combination of cytokinins (6-benzyl amino purine - BAP = 2.0 mgL<sup>-1</sup> and Kinetin = 0.25 mgL<sup>-1</sup>) and auxins (IAA - Indole acetic acid = 1.0 mgL<sup>-1</sup>) was found to be best suited for initiating callus in leaf explants (Plate 1) and used for leaf explants harvested from all varieties of green gram (Plate 2).

### In vitro callus induction

In leaf explants proliferation of callus occurred only in two green gram varieties out of the three varieties taken for study (Table 1; Plate 2, 3). Callus induction was observed in control leaf explants of CO-8 and both in control leaf explants as well as in leaf explants harvested from *in situ* supplementary UV-B irradiated NVL-585 crops. However, all leaf explants from VAMBAN-2 failed to induct callus. The induction of callus was delayed by one day in explants harvested from *in situ* UV-B irradiated NVL-585 compared with the control (Table 1). However, in CO-8 callus was induced one day earlier to NVL-585 control. The callus of *in situ* UV-B stressed NVL-585 variety of green gram weighed less by 51.28 % compared to control. The control CO-8 had better callus as evident by an accumulation in the fresh biomass by 10 % over the control callus of NVL-585. The trend observed in fresh weight continued in dry weight of callus also. The callus of *in situ* UV-B irradiated NVL-585 variety weighed less by 78.42 % below control on 30 DAI, while CO-8 control recorded the maximum (Table 1). The parenchyma cells of calluses proliferated from leaf explants were isodiametric with thin cell walls and were distributed evenly all through the callus in control samples (Plate 4, Fig. 1, 2). The parenchyma cells that have proliferated from the *in situ* UV-B irradiated callus were smaller by 29.67 % and more in number by 17 % in NVL-585 than their control (Plate 4, Fig. 2, 3). This is in accordance with the findings of Rajendiran *et al.* (2014a) who have reported the failure of leaf explants of some of the varieties of cowpea to proliferate callus after ultraviolet-B irradiation under *in vitro* culture. Rajendiran *et al.* (2014b) and Rajendiran *et al.* (2014c) opine that the response shown by stem explants and seeds to *in vitro* culture differ based on the sensitivity of the crops to *in vitro* culture conditions. Rajendiran *et al.* (2015a) in *Amaranthus dubius* Mart. Ex. Thell., Rajendiran *et al.* (2015e) in *Macrotyloma uniflorum* (Lam.) Verdc., Rajendiran *et al.* (2015f) in *Momordica charantia* L., Rajendiran *et al.* (2015g) in *Spinacia oleracea* L., Rajendiran *et al.* (2015h) in *Trigonella foenum-graecum* (L.) Ser., Rajendiran *et al.* (2015i) in *Benincasa hispida* (Thunb.) Cogn. and Rajendiran *et al.* (2015j) in *Portulaca oleracea* L. have also recorded varied results with leaf explants of *in situ* UV-B stressed plants under *in vitro* culture. In addition, *in vitro* germination of F<sub>1</sub> seeds harvested from *in situ* UV-B stressed black gram (Thiruvarasan and Rajendiran 2015), cowpea (Gowsalya and Rajendiran 2015) and green gram varieties (Vidya and Rajendiran 2015) also showed differences in viability.

Fig. 1: K = 0.1 mgL<sup>-1</sup>Fig. 2: K = 0.25 mgL<sup>-1</sup>Fig. 3: K = 0.5 mgL<sup>-1</sup>

**Plate 1:** Standardisation of Kinetin (K) concentration in culture media for *in vitro* regeneration from leaf explants using *Vigna radiata* (L.) Wilczek var. NVL-585 control samples. (7 DAI - Days after inoculation)

**Table 1:** Characteristics of callus proliferation in leaf explants of three varieties of 30 DAI *Vigna radiata* (L.) Wilczek from control and supplementary UV-B exposed conditions - *In vitro*.

Varieties	Treatment	Time taken for initiation (d)	Fresh weight (g)	Dry weight (g)	Parenchyma cell Frequency (µm)	Parenchyma cell size (µm)	
						Length	Breadth
CO-8	Control	23	0.896	0.157	479. ±0.45	147.57±0.83	142.54±1.76
	UV-B	-	-	-	-	-	-
NVL-585	Control	24	1.564	0.576	389.47±1.43	124.66±0.17	78.54±0.76
	UV-B	25	0.762	0.124	323.77±1.45	87.67±0.45	65.67±0.76
VAMBAN-2	Control	-	-	-	-	-	-
	UV-B	-	-	-	-	-	-

**Table 2:** The similarity indices in callus proliferation from leaf explants of three varieties of *Vigna radiata* (L.) Wilczek after supplementary UV-B exposure - *In vitro*.

Varieties	CO-8	NVL-585	VAMBAN-2
CO-8	100%	42.86%	14.29%
NVL-585	42.86%	100%	-
VAMBAN-2	14.29%	-	100%



Fig. 1: CO-8

Control



UV-B



Fig. 2: NVL-585

Control



UV-B



Fig. 3: VAMBAN-2

Control



UV-B

**Plate 2:** Comparison of *in vitro* callus proliferation from leaf explants of three varieties of *Vigna radiata* (L.) Wilczek on 30 DAI (Days after inoculation).



Fig. 1: CO-8 - Control



Fig. 2: NVL-585 - Control



Fig. 3: NVL-585 - UV-B

**Plate 3:** A closer view of calluses formed in two out of three varieties of *Vigna radiata* (L.) Wilczek from leaf explants of control and UV-B irradiated plants.




Fig. 1: CO-8 - Control




Fig. 2: NVL-585 - Control





Fig. 3: NVL-585 - UV-B

**Plate 4:** Cross section of calluses formed in two out of three varieties of *Vigna radiata* (L.) Wilczek from leaf explants of control and UV-B irradiated plants. (All figs. 400x)

#### **Dendrogram**

The variation shown by leaf explants of three varieties of green gram after *in situ* supplementary UV-B radiation in the parameters *viz.*, time taken for callus initiation, fresh and dry weight of callus, frequency and size of parenchyma cells in callus was reflected in the dendrogram. The two varieties *viz.*, CO-8 and NVL-585

had 42.86 % similarity and they formed one group as their leaf explants harvested from either control or UV-B stressed crops or both proliferated callus. VAMBAN-2 remained alone in the cluster showing very little similarity (14.29 %) with other two varieties, as both control and UV-B stressed explants failed to induct callus (Table 2; Plate 5).



**Plate 5:** Dendrogram showing the interrelationship between the three varieties of *Vigna radiata* (L.) Wilczek in callus proliferation from leaf discs of control and supplementary UV-B irradiated plants - *In vitro*.

## Conclusion

From the present experiment it is evident that the leaves of NVL-585 variety of green gram are the appropriate explants for germplasm conservation for surviving in UV-B elevated environment, as they are the only plant material to respond to *in vitro* culture after UV-B irradiation.

## Acknowledgement

The authors thank Prof. Dr. Thamizharasi Tamizhmani, Director, KMCPGS, Puducherry for providing research facilities and Dr. M. P. Ramanujam for his support and encouragement.

## References

- Arulmozhi D and Rajendiran K (2014a) Effect of supplementary ultraviolet-B radiation on nodulation and nitrogen metabolism in *Lablab purpureus* L. var. Goldy. *International Journal of Advanced Biological Research*. **4(3)**: 343-346.
- Arulmozhi D and Rajendiran K (2014b) Supplementary ultraviolet-B induced reduction in nodulation and nitrogen metabolism in hyacinth bean. *International Journal of Geology, Earth and Environmental Sciences*. **4(2)**: 73-77.
- Arulmozhi D and Rajendiran K (2014c) Effect of elevated ultraviolet-B irradiation on the nodulation and nitrogen metabolism in *Vigna unguiculata* (L.) Walp. cv. COFC-8. *International Journal of Food, Agriculture and Veterinary Sciences*. **4(2)**: 184-188.
- Bhat TM and Kudesia R (2011) Evaluation of Genetic Diversity in Five Different Species of Family Solanaceae using Cytological Characters and Protein Profiling. *Genetic Engineering and Biotechnology Journal*. 1- 8.
- Caldwell MM, Bjorn LO, Bornman JF, Flint SD, Kulandaivelu G, Teramura AH, Tevini M (1998) Effects of increased solar ultraviolet radiation on terrestrial ecosystems. *Photochem. Photobiol.* **46**: 40 - 52. DOI: 10.1016/S1011-1344(98)00184-5
- Everst B (1985) Clustering analysis, John Wiley and Sons, New York.
- Gowsalya L and Rajendiran K (2015) *In vitro* germination of F<sub>1</sub> seeds of cowpea varieties cultivated under *in situ* UV-B radiation. *International Journal of Biotechnology*. **4(4)**: 14-21.
- Kokilavani V and Rajendiran K (2013) Ultraviolet-B induced changes in the leaf epidermal and anatomical characteristics of *Vigna mungo* L. var. KM-2. *International Journal of Science and Nature*. **5(1)**: 126-130.
- Kokilavani V and Rajendiran K (2014a) Changes in leaf architecture of *Vigna unguiculata* (L.) Walp. cv. BCP-25 after exposure to elevated ultraviolet-B radiation. *International Journal of Science and Nature*. **5(3)**: 542-546.
- Kokilavani V and Rajendiran K (2014b) Ultraviolet-B induced changes in the leaf architecture of *Cucumis sativus* L. var. CO 1. *International Journal of Geology, Earth and Environmental Sciences*. **4(2)**: 208-215.
- Kokilavani V and Rajendiran K (2014c) Alterations in leaf architecture of *Ocimum sanctum* L. under elevated ultraviolet-B stress. *Global Journal of Bio-Science and Biotechnology*. **3(4)**: 374-378.
- Kokilavani V and Rajendiran K (2014d) Ultraviolet-B induced changes in the leaf epidermal and anatomical characteristics of *Vigna mungo* L. var. KM-2. *International Journal of Advanced Biological Research*. **5(1)**: 126-130.
- Kokilavani V and Rajendiran K (2014e) Effect of supplementary UV-B radiation on the yield of ten varieties of cowpea. *International Journal of Geology, Earth and Environmental Sciences*. **4(3)**: 65-73.
- Kokilavani V and Rajendiran K (2014g) Evaluation of the impact of Ultraviolet-B radiation on the foliar epidermal and anatomical characteristics of *Vigna unguiculata* (L.) Walp. c.v. COVU-1. *International Journal of Innovative Research and Review*. **2(4)**: 61-68.
- Kokilavani V and Rajendiran K (2014h) Variation in leaf architecture of *Vigna unguiculata* (L.) Walp. c.v. COFC-8 induced by supplementary UV-B exposure. *International Journal of Innovative Research and Review*. **2(4)**: 69-76.
- Kokilavani V and Rajendiran K (2014i) Ultraviolet-B induced reduction in nodulation in ten varieties of cowpea. *International Journal of Innovative Research and Review*. **2 (4)**: 77-82.
- Kokilavani V and Rajendiran K (2014j) Efficacy of *Vigna unguiculata* (L.) Walp. cv. Vamban leaves to withstand supplementary ultraviolet-B irradiation. *International Journal of Geology, Earth and Environmental Sciences*. **4(3)**: 203-210.
- Kokilavani V and Rajendiran K (2014k) Anatomical and epidermal alterations in the leaves of *Vigna unguiculata* (L.) Walp. cv. CO-6 due to UV-B exposure. *International Journal of Geology, Earth and Environmental Sciences*. **4(3)**: 211-218.

- Kokilavani V and Rajendiran K (2014l) A survey on the adaptive mechanism in leaf architecture of *Vigna unguiculata* (L.) Walp. cv. KM-1 under ultraviolet-B radiation. *International Journal of Food, Agriculture and Veterinary Sciences*. **4(3)**: 50-57.
- Kokilavani V and Rajendiran K (2014m) Modifications in leaf architecture of *Vigna unguiculata* (L.) Walp. cv. COVU-2 to defend from ultraviolet-B radiation. *International Journal of Food, Agriculture and Veterinary Sciences*. **4(3)**: 65-72.
- Kokilavani V and Rajendiran K (2014n) Analysis of the UV-B induced changes in morphology, anatomy and epidermis of *Vigna unguiculata* (L.) Walp. cv. CO-1 leaves. *International Journal of Food, Agriculture and Veterinary Sciences*. **4(3)**: 87-94.
- Kokilavani V and Rajendiran K (2014o) Influence of elevated ultraviolet-B radiation on the morphology and growth of ten varieties of cowpea. *International Journal of Food, Agriculture and Veterinary Sciences*. **4(3)**: 171-189.
- Kokilavani V and Rajendiran K (2015a) Variations in foliar morphology and anatomy of *Vigna unguiculata* (L.) Walp. c.v. CO-3 after supplementary ultraviolet-B exposure. *International Journal of Advanced Biological Research*. **5(1)**: 23-28.
- Kokilavani V and Rajendiran K (2015b) Study of leaf architecture of *Vigna unguiculata* (L.) Walp. cv. Pudukai under elevated ultraviolet-B radiation. *International Journal of Advanced Biological Research*. **5(1)**: 34-39.
- Kokilavani V and Rajendiran K (2014f) Influence of elevated Ultraviolet-B radiation on foliar organisation in *Vigna unguiculata* (L.) Walp. c.v. CW-122. *International Journal of Innovative Research and Review*. **2(4)**: 53-60.
- Kulandaivelu G, Maragatham S, Nedunchezian N (1989) On the possible control of ultraviolet - B induced response in growth and photosynthetic activities in higher plants. *Physiol. Plant*. **76**: 398 - 404. DOI: 10.1111/j.1399-3054.1989.tb06210.x
- Mark SM and Tevini M (1997) Effects of solar UV-B radiation on growth, flowering and yield of central and southern European bush bean cultivars (*Phaseolus vulgaris* L.). *Plant Ecology* **128**: 114-125. DOI: 10.1023/A:1009798528605
- Maskay N (1998) Single linkage clustering. In: Armitage P and Cotton T (Eds) *Encyclopedia of Biostatistics*, **Chapter 5**, Wiley, New York, 4121- 4122.
- Murashige T and Skoog F (1962) A revised medium for rapid growth and bioassays with tobacco tissue cultures. *Physiol Plant*. **15**: 473 - 497. DOI: 10.1111/j.1399-3054.1962.tb08052.x
- Rajendiran K (2001) Amelioration of Ultraviolet-B radiation impacts in green gram by Triadimefon. PhD. Thesis, Pondicherry University.
- Rajendiran K and Ramanujam MP (2003) Alleviation of ultraviolet-B radiation-induced growth inhibition of green gram by triadimefon. *Biologia Plantarum*. **46**: 621-624. DOI: 10.1023/A:1024840301092
- Rajendiran K and Ramanujam MP (2004) Improvement of biomass partitioning, flowering and yield by triadimefon in UV-B stressed *Vigna radiata* (L.) Wilczek. *Biologia Plantarum*. **48**: 145-148. DOI: 10.1023/B:BIOP.0000024293.31266.a2
- Rajendiran K, Ariswary D, Arulmozhi D and Kokilavani V (2015a) Screening of explants of ultraviolet-B exposed *Amaranthus dubius* Mart. Ex. Thell. for *in vitro* propagation. *International Journal of Food, Agriculture and Veterinary Sciences*. **5(2)**: 33-45.
- Rajendiran K, Gowsalya L and Sudaroli Sudha J (2015b) Assessment of variations in morphology and growth of three varieties of cowpea under elevated ultraviolet-B radiation. *International Journal of Food, Agriculture and Veterinary Sciences*. **5(30)**: 80-94.
- Rajendiran K, Gowsalya L, Vidya S and Thiruvarasan K (2015c) Influence of elevated ultraviolet-B radiation on the cotyledonary epidermis of F<sub>1</sub> seedlings of *Lagenaria siceraria* (Molina) Standl. var. Warad. *International Journal of Food, Agriculture and Veterinary Sciences*. **5(2)**: 98-103.
- Rajendiran K, Gowsalya L, Vidya S and Thiruvarasan K (2015d) Modifications in the cotyledonary epidermis of F<sub>1</sub> seedlings of *Lablab purpureus* (L.) Sweet var. Ankur under UV-B stress. *International Journal of Geology, Earth and Environmental Sciences*. **5(2)**: 188-193.
- Rajendiran K, Iswarya R, Arulmozhi D and Kokilavani V (2015e) Evaluation of ultraviolet-B stressed explants of *Macrotyloma uniflorum* (Lam.) Verdc. For *in vitro* propagation. *International Journal of Innovative Research and Review*. **3(2)**: 19-31.
- Rajendiran K, Julie Soniya F, Vijayalakshmi R and Kokilavani V (2015f) *In vitro* propagation of ultraviolet-B stressed *Momordica charantia* L. *International Journal of Geology, Earth and Environmental Sciences*. **5(2)**: 92-104.
- Rajendiran K, Kokilavani V and Murugananthan P (2014a) *In vitro* callus proliferation from leaf explants of ten varieties of cowpea after *in situ* ultraviolet-B irradiation. *International Journal of Geology, Earth and Environmental Sciences*. **4(3)**: 268-277.
- Rajendiran K, Kokilavani V and Murugananthan P (2014b) *In vitro* callus proliferation from stem explants of ten varieties of cowpea after *in situ* ultraviolet-B exposure. *International Journal of Food, Agriculture and Veterinary Sciences*. **4 (3)**: 190-199.
- Rajendiran K, Kokilavani V and Murugananthan P (2014c) *In vitro* seed germination and growth of ten varieties of cowpea after ultraviolet-B irradiation. *International Journal of Food, Agriculture and Veterinary Sciences*. **4(3)**: 200-217.
- Rajendiran K, Periyalakshmi D, Vijayalakshmi R and Kokilavani V (2015g) *In vitro* propagation of explants from

- ultraviolet-B stressed *Spinacia oleracea* L. *International Journal of Innovative Research and Review*. **3(2)**: 32-44.
- Rajendiran K, Priyadarsini V, Sudaroli Sudha J and Kokilavani V (2015h) *In vitro* propagation of *Trigonella foenum-graecum* (L.) Ser. after supplementary UV-B irradiation. *International Journal of Food, Agriculture and Veterinary Sciences*. **5(2)**: 46-57.
- Rajendiran K, Ramanujam MP (2006) Interactive effects of UV-B irradiation and triadimefon on nodulation and nitrogen metabolism in *Vigna radiata* plants. *Biologia Plantarum*. **50(4)**: 709-712. DOI: 10.1007/s10535-006-0112-3
- Rajendiran K, Riswan A, Sudaroli Sudha J and Kokilavani V (2015i) *In vitro* propagation of *Benincasa hispida* (Thunb.) Cogn. explants after enhanced UV-B exposure. *International Journal of Innovative Research and Review*. **3(2)**: 45-56.
- Rajendiran K, Shanmathy M, Sudaroli Sudha J and Kokilavani V (2015j) Assessment of ultraviolet-B tolerance in *Portulaca oleracea* L. for *in vitro* propagation. *International Journal of Biotechnology*. **4(2)**: 32-42.
- Rajendiran K, Thiruvarasan K and Vijayalakshmi R (2015k) Influence of elevated ultraviolet-B radiation on the morphology and growth of three varieties of black gram. *International Journal of Geology, Earth and Environmental Sciences*. **5(3)**: 62-76.
- Rajendiran K, Thiruvarasan K and Vijayalakshmi R (2015l) Assessment of nodulation in three varieties of black gram under elevated UV-B radiation. *International Journal of Geology, Earth and Environmental Sciences*. **5(3)**: 77-81.
- Rajendiran K, Thiruvarasan K and Vijayalakshmi R (2015m) Yield attributes of three varieties of black gram under *in situ* supplementary UV-B irradiation. *International Journal of Food, Agriculture and Veterinary Sciences*. **5(3)**: 67-74.
- Rajendiran K, Vidya and Arulmozhi D (2015n) Impact of supplementary UV-B radiation on the morphology and growth of *in situ* grown three green gram varieties. *International Journal of Geology, Earth and Environmental Sciences*. **5(3)**: 82-97.
- Rajendiran K, Vidya S and Arulmozhi D (2015o) Impact of supplementary ultraviolet-B radiation on the nodulation in three varieties of green gram. *International Journal of Food, Agriculture and Veterinary Sciences*. **5(3)**: 75-79.
- Rajendiran K, Vidya S, Gowsalya L and Thiruvarasan K (2015p) Impact of supplementary UV-B radiation on the morphology, growth and yield of *Vigna mungo* (L.) Hepper var. ADT-3. *International Journal of Food, Agriculture and Veterinary Sciences*. **5(2)**: 104-112.
- Rajendiran K, Vidya S, Gowsalya L and Thiruvarasan K (2015q) Changes in the morphology, growth and yield of *Vigna mungo* (L.) Hepper var. VAMBAN-4 under ultraviolet-B stress. *International Journal of Food, Agriculture and Veterinary Sciences*. **5(3)**: 59-66.
- Sudaroli Sudha J and Rajendiran K (2013a) Effect of elevated UV-B irradiation on the nodulation and nitrogen metabolism in *Sesbania grandiflora* (L.) Pers. *International Journal of Science and Nature*. **4(4)**: 664-667.
- Sudaroli Sudha J and Rajendiran K (2013b) Effect of elevated UV-B irradiation on the nodulation and nitrogen metabolism in *Vigna unguiculata* (L.) Walp. c.v. BCP-25. *International Journal of Food, Agriculture and Veterinary Sciences*. **3(3)**: 77-81.
- Sudaroli Sudha J and Rajendiran K (2014a) Impact of ultraviolet-B radiation on nodulation and nitrogen metabolism in *Vigna unguiculata* (L.) Walp. cv. COVU-1. *International Journal of Geology, Earth and Environmental Sciences*. **4(2)**: 224-230.
- Sudaroli Sudha J and Rajendiran K (2014b) Ultraviolet-B induced reduction in nodulation and nitrogen metabolism in *Vigna mungo* (L.) Hepper var. T-9. *Global Journal of Bioscience and Biotechnology*. **3(4)**: 370-373.
- Sudaroli Sudha J, Rajendiran K (2014c) Ultraviolet-B induced reduction in nodulation and nitrogen metabolism in *Vigna unguiculata* (L.) Walp. cv. CO-1. *Global Journal of Bioscience and Biotechnology*. **4(3)**: 10-14.
- Sullivan JH, Teramura AH and Dillenburg LR (1994) Growth and photosynthetic responses of field-grown sweetgum (*Liquidambar styraciflua*) seedlings to UV-B radiation. *American Journal of Botany*. **81**: 826-832. DOI: 10.2307/2445763
- Thiruvarasan K and Rajendiran K (2015) Screening of F<sub>1</sub> seeds of black gram varieties grown under *in situ* UV-B radiation for *in vitro* germination. *International Journal of Biotechnology*. **4(4)**: 22-29.
- Vidya S and Rajendiran K (2015) *In vitro* germination of F<sub>1</sub> seeds harvested from *in situ* UV-B stressed green gram varieties. *International Journal of Biotechnology*. **4(4)**: 6-13.
- Vijayalakshmi R and Rajendiran K (2014a) Impact of ultraviolet-B radiation on nodulation and nitrogen metabolism in *Cyamopsis tetragonoloba* (L.) Taub. var. PNB. *International Journal of Geology, Earth and Environmental Sciences*. **4(2)**: 78-82.
- Vijayalakshmi R and Rajendiran K (2014b) Impact of ultraviolet-B radiation on nodulation and nitrogen metabolism in *Phaseolus vulgaris* L. cv. Prevail. *International Journal of Advanced Biological Research*. **4(3)**: 339-342.
- Vijayalakshmi R and Rajendiran K (2014c) Effect of elevated ultraviolet-B irradiation on the nodulation and nitrogen metabolism in *Vigna unguiculata* (L.) Walp. cv. CW-122. *International Journal of Food, Agriculture and Veterinary Sciences*. **4(2)**: 189-193.